




Mississippi Valley Conservation

*State of the Lake
Environment Report
December 2007*

Joes Lake



Joes Lake

Joes Lake is situated in the Township of Lanark Highlands, in Lanark County. It is one of the lakes of the Clyde River system, in the Mississippi Valley watershed. Joes Lake is at an elevation of 195 metres above sea level. The lake perimeter is approximately 5.3 kilometres, with a maximum depth of 4.3 metres. Joes Lake supports a warm water fishery. Common species include Yellow Pickerel and Northern Pike. At last count in 1983, there were 19 cottages and one campsite (30 sites) on the lake.



Limited water quality data is available for Joes Lake. Records indicate that shoreline property owners have not yet formed a Lake Association or participated in the Ministry of Environment's Self-Help or Lake Partner Program. Comprehensive testing in 2002 and 2007 through Mississippi Valley Conservation's (MVC) *Watershed Watch Program* provides for a comparison between water quality conditions as they exist now, to results obtained 24 years ago through the MOE Recreational Lakes Program.

In general, the water quality in Joes Lake has improved. There is one sampling station at the deepest point, in the west end of the lake. This station was sampled three times in 2007. Graphs will follow that show water clarity, as measured by secchi disc. The average reading for 2007 was 3.7 metres, indicating Joes Lake is a moderately enriched (some nutrients) or mesotrophic lake. Twenty four years ago, the average secchi disc depth was 3.0 metres, indicating a slight improvement in water clarity.

Directly related to water clarity is the amount of nutrients, in particular phosphorus, entering the lake. The Provincial Objective for phosphorus levels in shield lakes is a maximum of 20 micrograms per litre (ug/L). In 2007, the average total phosphorus level in Joes Lake was 12.0 ug/L , indicating that the lake environment is moderately enriched (some nutrients) or mesotrophic. Twenty four years ago, the average phosphorus level was 22.0 ug/L , over the Provincial Objective. Joes Lake decreased its average phosphorus levels by 10.0 ug/L , changing its trophic status from a Eutrophic lake environment (enriched, high levels of nutrients) to a moderately enriched lake (some nutrients) and putting it well below the Provincial Objective.

Chlorophyll-a, is a measure of the algal density in the lake. The average chlorophyll-a density for the sampling station was 3.33 ug/L , indicating a moderate algal density for Joes Lake in 2007. In 1983, chlorophyll-a levels were slightly lower at 3.15 ug/L .

Plants and animals are a direct reflection of their environment. The most critical time of year for conducting dissolved oxygen and temperature profiles is after August 31. Profiles are generally conducted at this time of year and at the deepest point in the lake. Aquatic vegetation and algae that has grown over the summer, has died off and settled on the bottom, using the available oxygen necessary to sustain aquatic life in the lower portion of the lake or the hypolimnion. Two other profiles were conducted in 2007, one in May and one in July, in order to generate a more concise picture of the oxygen content of the lake.

The dissolved oxygen and temperature data, measured at the deepest point in the main basin, indicate oxygen concentrations in the deep water portion have decreased by late summer. Warm water fish species, such as pike and pickerel, are squeezed into the upper two metres of the lake by September. Therefore, residents and users of Joes Lake cannot afford to be complacent. Every effort should be made to reduce nutrient loading into the lake from land use activities.

Joes Lake was also tested for invasive species in 2007, in particular, for zebra mussels and spiny water flea, in partnership with the Ontario Federation of Anglers and Hunters. Joes Lake did *not* have zebra mussel veligers (larvae) or spiny water flea present in the samples collected. Residents and property owners need to ensure that all access points to the lake have posted signs indicating the precautions they can take to avoid the spread of invasive species into Joes Lake. Another recommendation is for residents to begin participation in the invasive species monitoring program through MVC.



Residents and users of Joes Lake need to adopt a stewardship approach to limit the amount of nutrients entering the lake. The first step to achieve this, is to form a Lake Association. It is recommended that a Lake Steward be appointed to undertake ongoing water quality testing and to join the Mississippi Valley Lake Stewardship Network. Monitoring over time is essential to determine long term trends and changes. Resources and information are readily available through the *Watershed Watch Program*. There are helpful tips throughout this report to help reduce your impact on Joes Lake. Additional water quality data, current and historic, is available for Joes Lake and many other lakes in the Mississippi Valley watershed. Contact MVC for more information on how you can become a good lake steward for your lake. We all have a responsibility to preserve this precious natural resource for future generations.

FIVE EASY STEPS TO IMPROVE WATER QUALITY



1. Build at least 30 metres away from the shoreline.
2. Keep your lot well treed and preserve or replant native vegetation along the shoreline.
3. Pump out your septic tank every three to five years.
4. Reduce water use and use phosphate free soaps and detergents.
5. Keep the size of your lawn to a minimum; do not use fertilizers, herbicides or pesticides.

LOW PHOSPHORUS LIFESTYLE	Amount of Phosphorus (grams)	HIGH PHOSPHORUS LIFESTYLE	Amount of Phosphorus (grams)
Human waste	535	Human waste	535
No dishwasher	0	Dishwasher using powdered detergent once per day	650
No fertilizer	0	Lawn fertilized once/year	1960
Trees not cut down	20	Lot cleared of trees	30
Uses phosphate-free products	20	Uses products with phosphate	180
TOTAL	575 grams	TOTAL	3355 grams

Joes Lake




A Secchi Disc visually measures water clarity. The depth at

which the disc disappears indicates the level of nutrients and algae growth. The higher the reading, the clearer the lake. The more nutrients that run into the lake, the more algae growth, thus causing reduced water clarity.

Secchi disc readings taken in 2007 indicate good clarity with the mean being 3.7 m, indicating a moderately enriched or mesotrophic lake.

This lake was last surveyed using imperial measurements, therefore the original bathymetric (depth) measurements are recorded in feet rather than metres.

3.28 ft = 1 m 1ft = 0.3048 m



Most Common Fish Species
 Yellow Pickerel
 Northern Pike

Latitude	45° 05'
Longitude	76° 10'
Max. Depth	4.3 m
Mean Depth	3.7 m
Perimeter	5.3 km
Surface Area	59.94 ha
Volume	6.36 m ³ x 10 ⁷
Height above sea level	195 m

↑ Watershed Watch
 Sampling Station

This map is intended for illustration only; it should not be used as a navigation guide.

*Remember
 Use non-lead sinkers to protect the
 health of the fish and this lake.*

How Does Joes Lake Measure Up?

1983-2007 Water Quality Results

Sample Year Mean	Secchi Disc Depth [Metres]	Total Phosphorus Euphotic Zone [Micrograms/litre]	Chlorophyll-a Composite [Micrograms/Litre]
**1983	3.0	23.2	3.15
2002	3.3	14.8	7.64
2007	3.7	12.0	3.33
n	3	3	3
Minimum	3.0	12.0	3.15
Maximum	3.7	23.2	7.64
Mean	3.3	16.6	4.70
Standard Deviation	3.4	5.8	2.54

*Mean based on less than 6 measurements **Includes Recreational Lakes Program Data
 Chlorophyll-a data prior to 1985 has been adjusted to reflect new lab procedures filtering resulting in an increase in chla concentrations by 35%

Interpreting Secchi Disc Readings:


A Secchi disc is a black and white coloured disc used to determine water clarity. The disc is lowered into the water. The point, at which you can no longer distinguish the black and white, is called the secchi depth.



The higher the Secchi Disc measurement the clearer your lake is.

INTERPRETING YOUR SECCHI DISC RESULTS	
Secchi Reading	Lake Nutrient Status
Over 5 metres	Oligotrophic - unenriched, few nutrients
3.0 to 4.9 metres	Mesotrophic – moderately enriched, some nutrients
Less than 2.9 metres	Eutrophic – enriched, higher levels of nutrients


Joe's Lake Annual Secchi Disc Results



Interpreting Total Phosphorus Results:

Phosphorus is the nutrient that controls the growth of algae in most Ontario lakes. For this reason any increase in phosphorus in the lake will increase the quantity of algae that can grow. High levels of phosphorus can lead to algal blooms and in some cases affect the habitat of cold water fish such as lake trout. A general guideline exists to characterize your lake based on the total phosphorus that is measured.

INTERPRETING YOUR TOTAL PHOSPHORUS RESULTS	
Total Phosphorus	Lake Nutrient Status
10ug/L or less	Oligotrophic - unenriched, few nutrients
11 to 20ug/L	Mesotrophic - moderately enriched, some nutrients
21ug/L or more	Eutrophic - enriched, higher levels of nutrients



How to protect or restore a shoreline depends on the conditions of the site and the energy and resources of the owner.

There are four main strategies to choose from:

PRESERVATION

When purchasing a lakefront property, a natural shoreline is retained and access to the lake is designed to avoid shoreline damage.

ENHANCEMENT

Native species are planted and non-native species are removed.

NATURALIZATION

Degraded shorelines are left alone to return to their natural state.

RESTORATION


Cleared areas are planted with native species.




Interpreting Chlorophyll-a Results:

Chlorophyll-a, is a measure of the algal density in the lake. The lower the chlorophyll-a density in your lake, the clearer your lake is. Chlorophyll-a, is directly affected by the amount of total phosphorus in your lake. The more phosphorus there is in the water, the more algal growth will occur.

INTERPRETING YOUR CHLOROPHYLL-A RESULTS	
Chlorophyll-a Reading	Lake Nutrient Status
Up to 2 ug/L - low algal density	Oligotrophic - unenriched, few nutrients
2-4 ug/L - moderate algal density	Mesotrophic - moderately enriched, some nutrients
More than 4 ug/L - high algal density	Eutrophic - enriched, higher levels of nutrients




**For more information on lakes in the
Mississippi Valley Watershed, visit MVC
online at
www.mvc.on.ca**



Interpreting pH Results:


The pH value is a measure of the concentration of hydrogen ions of a substance, which ranges from very acidic (pH = 1) to very alkaline (pH = 14). At a normal to neutral acidity level, a lake supports a diversity of life. A pH of 7 is neutral and most lake waters range between 6 and 9. pH values less than 6 are considered acidic, and most life forms cannot survive at a pH of 4.0 or lower. This parameter directly influences the types of plants and animals that live in the lake. Lakes with pH levels at 7.3 or higher are vulnerable to zebra mussels invasive.



φβφβφβφβφβφβφβφβφβφβφβφβφ

MVC and O.F.A.H. need your help to Stop the Invasion!

Check & clean your boat every time you change water bodies



Working with Lake Associations, we hope to improve signage at public launching areas to identify lakes where zebra mussels and spiny water fleas are already present. We hope to focus on an ambitious educational campaign to help reduce their spread to lakes where they are not yet present.

For more information call MVC at (613)259-2421, the Invading Species Hotline 1-800-563-7711.

φβφβφβφβφβφβφβφβφβφβφβφβφ

JOES LAKE – Main Basin DISSOLVED OXYGEN/TEMPERATURE PROFILES



Depth [Metres]	Temperature [Degrees Celsius]	Dissolved Oxygen [Milligrams/Litre]	Percent % Saturation	Stratification
0.1	14.0	10.0	95	Epilimnion
1.0	14.0	10.6	99	
2.0	13.9	11.3	106	
3.0	12.8	11.4	101	Thermocline
4.0	12.3	11.5	105	Hypolimnion
5.0	Bottom	Bottom	Bottom	

MOE Rec. Lks. Station # 19-3430-764-01, MVC Station # 02-07

Date: May 4, 2007

Depth: 5.0 Metres

Euphotic Zone (Penetration of Light) = 5.0 Metres

Depth [Metres]	Temperature [Degrees Celsius]	Dissolved Oxygen [Milligrams/Litre]	Percent % Saturation	Thermal Stratification
0.1	23.1	10.2	115	Epilimnion
1.0	21.9	12.2	135	Thermocline - 1
2.0	21.5	12.3	135	Thermocline - 2
3.0	21.0	11.3	124	Hypolimnion
4.0	18.7	6.5	66	
5.0	Bottom	Bottom	Bottom	

Date: July 3, 2007

Depth: 5.0 Metres

Euphotic Zone (Penetration of Light) = 5.0 Metres

Date: September 10, 2007


Depth: 4.0 Metres

Euphotic Zone (Penetration of Light) = 4.0 Metres



The Watershed Watch program was made possible thanks to the generous support of the Ministry of Environment, Lake Associations, area Stewardship Councils, the Lake Stewardship Network and concerned citizens.

For more information regarding Watershed Watch or for advice on how you can help protect or enhance your lake environment, contact Susan Lee, Watershed Monitoring Supervisor, Mississippi Valley Conservation at (613) 259-2421 or slee@mvc.on.ca



Mississippi Valley Conservation

Depth [Metres]	Temperature [Degrees Celsius]	Dissolved Oxygen [Milligrams/Litre]	Percent % Saturation	Thermal Stratification
0.1	21.2	7.0	75	Epilimnion
1.0	20.9	5.7	60	
2.0	20.7	4.8	50	
3.0	20.7	3.8	40	
4.0	Bottom	Bottom	Bottom	

*** Warm Water Fisheries Habitat (Bass, Walleye, Pike and Perch) defined as Dissolved Oxygen Concentrations greater than 4 mg/L at Temp. less than 25°C**



